

# Atomically Tailored Nanomagnets and their use for Atomic-Level Spintronics

Wiesendanger, Roland<sup>1</sup>

<sup>1</sup>*University of Hamburg, Institute of Applied Physics and Interdisciplinary Nanoscience Center Hamburg, D-20355 Hamburg, Germany*

The developments of novel magnetic materials as well as spin-based electronics are hot topics of current research in nanoscale science. Both research fields profit tremendously from atomic-scale insight into magnetic properties and spin-dependent interactions at the atomic level. Based on the development of spin-polarized scanning tunneling microscopy (SP-STM) [1] we have recently established the novel method of single-atom magnetometry [2,3] which allows the measurement of magnetization curves and the determination of magnetic moments on an atom-by-atom basis. While the sensitivity level of single-atom magnetometry is below one Bohr magneton, it can easily be combined with the atomic-resolution imaging and manipulation capabilities of conventional STM, thereby offering a novel approach towards a rational material design based on the knowledge of the atomic-level properties and interactions within the solid state [4]. Moreover, an atom-by-atom design and realization of all-spin logic devices [5] has recently been demonstrated by our group based on the combined knowledge derived from surface physics, nanoscience, and magnetism. By using SP-STM we have recently discovered nanoskyrmion lattices of single atomic layers of transition metals on particular substrates exhibiting a large spin-orbit coupling. In this case, skyrmionic lattices can be stabilized by Dzyaloshinskii-Moriya interactions combined with the breaking of inversion symmetry at surfaces and interfaces [6]. Following this approach, the existence of skyrmions of ultimate small size, being stable even in zero field, has recently been demonstrated, offering great potential for future nanospintronic devices.

[1] R. Wiesendanger, *Rev. Mod. Phys.* 81, 1495 (2009). [2] F. Meier et al., *Science* 320, 82 (2008). [3] L. Zhou et al., *Nature Physics* 6, 187 (2010). [4] A. A. Khajetoorians et al., *Nature Physics* 8, 497 (2012). [5] A. A. Khajetoorians et al., *Science* 332, 1062 (2011). [6] S. Heinze et al., *Nature Physics* 7, 713 (2011).